
Markov Chains

ECE 341: Random Processes

Lecture #20

note: All lecture notes, homework sets, and solutions are posted on www.BisonAcademy.com

A and B play a match

Problem 1: Two teams, A and B, are playing a match

- A has a 70% chance of winning any given game
- The first team to win 3 games wins the match.

This is a binomial distribution

Problem 2: Two teams, A and B, are playing a match

- A has a 70% chance of winning any given game
- The first team to win **by** 3 games wins the match.

This is a *totally* different problem.


Problem #2 is an infinite sequence

- To solve, we need a different tool: Markov chains.
-

Markov Chain


A Markov chain is a discrete-time probability function where

- $X(k)$ is the state of the system at time k , and
- $X(k+1) = A X(k)$



Three people, A, B, and C, are playing ball. Every second they pass the ball at random:

- When A has the ball, he/she
 - Keeps the ball 50% of the time
 - Passes it to B 20% of the time, and
 - Passes it to C 30% of the time
- When B has the ball, he/she
 - Passes it to A 30% of the time
 - Keeps it 60% of the time, and
 - Passes it to C 10% of the time
- When C has the ball, he/she
 - Passes it to A 40% of the time, and
 - Passes it to B 60% of the time.



Assume at $t=0$, A has the ball.

- What is the probability that B will have the ball after k tosses?
 - After infinite tosses?
-

This lecture covers three different methods to analyze problems of this sort:

- Matrix multiplication
- Eigenvalues and Eigenvectors, and
- z-Transforms.

Solution #1: Matrix Multiplication

Let $X(k)$ be the probability that A, B, and C have the ball at time k :

$$x(k) = \begin{bmatrix} p(a) \\ p(b) \\ p(c) \end{bmatrix}$$

Then from the problem statement, $X(k+1)$ is related to $X(k)$ by a state-transition matrix:

$$x(k+1) = \begin{bmatrix} 0.5 & 0.3 & 0.4 \\ 0.2 & 0.6 & 0.6 \\ 0.3 & 0.1 & 0 \end{bmatrix} x(k) = A x(k) \quad x(0) = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

Note that

- The columns are the probabilities in the above problem statement, and
 - The columns must add up to 1.000 (all probabilities add to one)
-

The probability of each player having the ball after 1, 2, 3 tosses is (using Matlab)

```
A = [0.5,0.2,0.3 ; 0.3,0.6,0.1 ; 0.4,0.6,0]'
```

```
    0.5000    0.3000    0.4000
    0.2000    0.6000    0.6000
    0.3000    0.1000         0
```

```
X = [1;0;0]      % k = 0
```

```
    1.0000
    0.0000
    0.0000
```

```
X = A*X          % k = 1
```

```
    0.5000
    0.2000
    0.3000
```

```
X = A*X          % k = 2
```

```
    0.4300
    0.4000
    0.1700
```

```
X = A*X           % k = 3
```

```
0.4030  
0.4280  
0.1690
```

time passes

```
X = A*X           % k = 100
```

```
0.3953  
0.4419  
0.1628
```

Eventually X quits changing. This is the steady-state solution.

Steady-State Solution:

If you want to find the steady-state solution, you can simply raise A to a large number (like 100) and solve in one shot:

$$X_0 = [1; 0; 0]$$

1
0
0

$$X_{20} = A^{100} * X_0$$

0.3953
0.4419
0.1628

You can also solve for the steady-state solution by finding $x(k)$ such that

$$x(k+1) = x(k) = A x(k)$$

Solving:

$$(A - I)x(k) = 0$$

$$\left(\begin{bmatrix} 0.5 & 0.3 & 0.4 \\ 0.2 & 0.6 & 0.6 \\ 0.3 & 0.1 & 0 \end{bmatrix} - \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} a \\ b \\ c \end{bmatrix} = 0$$

$$\begin{bmatrix} -0.5 & 0.3 & 0.4 \\ 0.2 & -0.4 & 0.6 \\ 0.3 & 0.1 & -1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = 0$$

Assume $c = 1$

$$\begin{bmatrix} -0.5 & 0.3 \\ 0.2 & -0.4 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = - \begin{bmatrix} 0.4 \\ 0.6 \end{bmatrix}$$

```
ab = -inv([-0.5, 0.3; 0.2, -0.4]) * [0.4; 0.6]
```

```
2.4286
```

```
2.7143
```

```
X = [ab; 1]
```

```
2.4286
```

```
2.7143
```

```
1.0000
```

```
X = X / sum(X)
```

```
0.3953
```

```
0.4419
```

```
0.1268
```

which is the same answer we got before.

Solution #2: Eigenvalues and Eigenvectors

The problem we're trying to solve is

$$x(k+1) = A x(k)$$

$$x(0) = X_0$$

This is actually an eigenvalue / eigenvector problem.

- Eigenvalues tell you how the system behaves,
 - Eigenvectors tell you what behaves that way.
-

Since this system has three states, the generalized solution for $x(k)$ will be:

$$x(k) = a_1 \Lambda_1 \lambda_1^k + a_2 \Lambda_2 \lambda_2^k + a_3 \Lambda_3 \lambda_3^k$$

where

- λ_i is the i th eigenvalue,
- Λ_i is the i th eigenvector, and
- a_i is a constant depending upon the initial condition.

At $k = 0$:

$$x(0) = \begin{bmatrix} \Lambda_1 & \Lambda_2 & \Lambda_3 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix}$$



The excitation of each eigenvector is then

$$x_0 = [1; 0; 0]$$

$$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

$$A_{123} = \text{inv}(M) * x_0$$

$$\begin{bmatrix} 0.6149 \\ 0.5482 \\ 0.9354 \end{bmatrix}$$

meaning

$$x(k) = 0.6149 \begin{bmatrix} 0.6430 \\ 0.7186 \\ 0.2468 \end{bmatrix} (1)^k + 0.5482 \begin{bmatrix} 0.2222 \\ 0.5693 \\ -0.7915 \end{bmatrix} (-0.1562)^k + 0.9543 \begin{bmatrix} 0.5151 \\ -0.8060 \\ 0.2989 \end{bmatrix} (0.2562)^k$$



or adding the scalars to the eigenvectors:

$$W = \text{inv}(M) * X0$$

```
0.6149
0.5482
0.9354
```

$$M * \text{diag}(W)$$

```
0.3953    0.1218    0.4828
0.4419    0.3121   -0.7539
0.1628   -0.4339    0.2711
```

$$x(k) = \begin{bmatrix} 0.3953 \\ 0.4419 \\ 0.1628 \end{bmatrix} (1)^k + \begin{bmatrix} 0.1218 \\ 0.3121 \\ -0.4339 \end{bmatrix} (-0.1562)^k + \begin{bmatrix} 0.4828 \\ -0.7539 \\ 0.2711 \end{bmatrix} (0.2562)^k$$

As k goes to infinity, the first eigenvector is all that remains.

Steady-State Solution using Eigenvectors

Note that the steady-state solution is simply the eigenvector associated with the eigenvalue of 1.000.

$$[M, V] = \text{eig}(A)$$

M = *eigenvectors*

0.6430	0.2222	0.5161
0.7186	0.5693	-0.8060
0.2648	-0.7915	0.2898

V = *eigenvalues*

1.0000	0	0
0	-0.1562	0
0	0	0.2562



Scale so the total is 1.0000

```
X = M(:, 1);  
X = X / sum(X)
```

```
0.3953
```

```
0.4419
```

```
0.1628
```

Same answer as before.

Solution #3: z-Transforms

Again, the problem we are trying to solve is

$$x(k+1) = \begin{bmatrix} 0.5 & 0.3 & 0.4 \\ 0.2 & 0.6 & 0.6 \\ 0.3 & 0.1 & 0 \end{bmatrix} x(k) \quad x(0) = X_0 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

This can be written as

$$x(k) = A x(k-1) + X_0 \delta(k)$$

$$x(k+1) = A x(k) + X_0 \delta(k+1)$$

Take the z-transform

$$zX = AX + zX_0$$

To determine the probability that B has the ball at time k, look at the second state

$$Y = CX = \begin{bmatrix} 0 & 1 & 0 \end{bmatrix} X$$

Solving for Y then gives the z-transform for b(k)

$$zX = AX + zX_0$$

$$(zI - A)X = zX_0$$

$$X = z(zI - A)^{-1} X_0$$

$$Y = CX = z C(zI - A)^{-1} X_0$$

$$Y = z C(zI - A)^{-1} X_0$$

For our 3x3 example,

$$Y(z) = z \begin{bmatrix} 0 & 1 & 0 \end{bmatrix} \left(\begin{bmatrix} z & 0 & 0 \\ 0 & z & 0 \\ 0 & 0 & z \end{bmatrix} - \begin{bmatrix} 0.5 & 0.3 & 0.4 \\ 0.2 & 0.6 & 0.6 \\ 0.3 & 0.1 & 0 \end{bmatrix} \right)^{-1} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

$\begin{matrix} \text{C} & \text{A} & \text{B} = \text{X0} \end{matrix}$

This is somewhat painful to compute by hand. Fortunately, there's Matlab to the rescue.

- $G = \text{ss}(A, B, C, D, T)$ *input a dynamic system into matlab*
 - $Y = \text{tf}(G)$ *calculate and express the z-transform of Y in transfer function form*
 - $Y = \text{zpk}(G)$ *calculate and express the z-transform of Y in factored form*
-

In matlab:

```
A = [0.5,0.3,0.4;0.2,0.6,0.6;0.3,0.1,0]
```

```
    0.5000    0.3000    0.4000  
    0.2000    0.6000    0.6000  
    0.3000    0.1000         0
```

```
X0 = [1;0;0]
```

```
    1  
    0  
    0
```

```
C = [0,1,0]
```

```
    0    1    0
```

```
Bz = ss(A, X0, C, 0, 1);
```

tf (Bz)

$$\frac{0.2 z + 0.18}{z^3 - 1.1 z^2 + 0.06 z + 0.04}$$

Sampling time (seconds): 1

zpk (Bz)

$$\frac{0.2 (z+0.9)}{(z-1) (z-0.2562) (z+0.1562)}$$

Sampling time (seconds): 1

(recall that you need to multiply by z)

Finding $b(k)$:

$$B(z) = \left(\frac{0.2(z+0.9)z}{(z-1)(z-0.2562)(z-0.1562)} \right)$$

factor out a z and use partial fractions:


$$B(z) = \left(\left(\frac{0.6054}{z-1} \right) + \left(\frac{-3.1089}{z-0.2562} \right) + \left(\frac{2.5034}{z-0.1562} \right) \right) z$$

multiply by z

$$B = \left(\frac{0.6054z}{z-1} \right) + \left(\frac{-3.1089z}{z-0.2562} \right) + \left(\frac{2.5034z}{z-0.1562} \right)$$

Take the inverse z -transform

$$b(k) = \left(0.6054 - 3.1089(0.2562)^k + 2.5034(0.1562)^k \right) u(k)$$



Probability that player B has the ball after toss k


z-Transform with Complex Poles

You can get complex poles. If you do, use entry in the z-transform table:

$$\left(\frac{(a\angle\theta)z}{z-b\angle\phi}\right) + \left(\frac{(a\angle-\theta)z}{z-b\angle-\phi}\right) \rightarrow 2a b^k \cos(\phi k - \theta) u(k)$$

For example, suppose player A, B, and C toss the ball as:

- A keeps the ball 30% of the time and passes it to B 70% of the time
- B keeps the ball 20% of the time and passes it to C 80% of the time, and
- C keeps the ball 10% of the time and passes it to A 90% of the time



Suppose A starts with the ball at $k = 0$. Determine the probability that B has the ball after k tosses.

Solving using z-transforms: express in matrix form

$$zX = \begin{bmatrix} 0.3 & 0 & 0.9 \\ 0.7 & 0.2 & 0 \\ 0 & 0.8 & 0.1 \end{bmatrix} X \quad X(0) = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

$$Y = p(B) = \begin{bmatrix} 0 & 1 & 0 \end{bmatrix} X$$



Find B(z) using Matlab

```
A = [0.3, 0, 0.9; 0.7, 0.2, 0; 0, 0.8, 0.1]
```

```
    0.3000         0    0.9000
    0.7000    0.2000         0
         0    0.8000    0.1000
```

```
X0 = [1; 0; 0];
```

```
C = [0, 1, 0];
```

```
Bz = ss(A, X0, C, 0, 1);
```

```
zpk(Bz)
```

```
    0.7 (z-0.1)
```

```
-----  
(z-1) (z^2 + 0.4z + 0.51)
```

Sampling time (seconds): 1

(again - multiply by z to get B(z))

$$B(z) = \left(\frac{0.7(z-0.1)z}{(z-1)(z-0.7142\angle 106^0)(z-0.7142\angle -106^0)} \right)$$

Pull out a z and expand using partial fractions


$$B(z) = \left(\left(\frac{0.3298}{(z-1)} \right) + \left(\frac{0.2764\angle -126.8^0}{(z-0.7142\angle 106^0)} \right) + \left(\frac{0.2764\angle 126.8^0}{(z-0.7142\angle -106^0)} \right) \right) z$$

Multiply both sides by z

$$B = \left(\frac{0.3298z}{(z-1)} \right) + \left(\frac{z0.2764\angle -126.8^0}{(z-0.7142\angle 106^0)} \right) + \left(\frac{z0.2764\angle 126.8^0}{(z-0.7142\angle -106^0)} \right)$$

Take the inverse z-transform

$$z b(k) = \left(0.3298 + 0.5527(0.7142)^k \cos(k \cdot 106^0 + 126.8^0) \right) u(k)$$



probability that player B has the ball after k tosses