


## Dynamics of a 2-Link Arm



Problem: Determine the dynamics of a 2-link arm with

- Each arm having a length of 1m, and
- Each arm has a lump-mass of 1kg at the end of each arm.

Step 1: Determine the (x,y) position and velocity of each mass

mass 1:

$$\begin{aligned} x_1 &= -\sin(\theta_1) & \dot{x}_1 &= -\cos(\theta_1)\dot{\theta}_1 \\ y_1 &= \cos(\theta_1) & \dot{y}_1 &= -\sin(\theta_1)\dot{\theta}_1 \end{aligned}$$

mass 2:

$$\begin{aligned} x_2 &= x_1 - \sin(\theta_1 + \theta_2) & \dot{x}_2 &= -\cos(\theta_1)\dot{\theta}_1 - \cos(\theta_1 + \theta_2)(\dot{\theta}_1 + \dot{\theta}_2) \\ y_2 &= y_1 + \cos(\theta_1 + \theta_2) & \dot{y}_2 &= -\sin(\theta_1)\dot{\theta}_1 - \sin(\theta_1 + \theta_2)(\dot{\theta}_1 + \dot{\theta}_2) \end{aligned}$$

Using shorthand notation

$$\begin{aligned} x_1 &= -s_1 & \dot{x}_1 &= -c_1\dot{\theta}_1 \\ y_1 &= c_1 & \dot{y}_1 &= -s_1\dot{\theta}_1 \end{aligned}$$

$$\begin{aligned} x_2 &= -s_1 - s_{12} & \dot{x}_2 &= -c_1\dot{\theta}_1 - c_{12}(\dot{\theta}_1 + \dot{\theta}_2) \\ y_2 &= c_1 + c_{12} & \dot{y}_2 &= -s_1\dot{\theta}_1 - s_{12}(\dot{\theta}_1 + \dot{\theta}_2) \end{aligned}$$

---

Step 2: Form the kinetic and potential energy:

Mass 1:

$$KE = \frac{1}{2}m(\dot{x}_1^2 + \dot{y}_1^2)$$

$$KE = \frac{1}{2}\left(\left(-c_1\dot{\theta}_1\right)^2 + \left(-s_1\dot{\theta}_1\right)^2\right)$$

Note that  $\sin^2 + \cos^2 = 1$

$$KE = \frac{1}{2}\dot{\theta}_1^2$$

$$PE = mgy_1$$

$$PE = gc_1$$

Mass 2:

$$KE = \frac{1}{2}m(\dot{x}_2^2 + \dot{y}_2^2)$$

$$KE = \frac{1}{2}\left(\left(-c_1\dot{\theta}_1 - c_{12}(\dot{\theta}_1 + \dot{\theta}_2)\right)^2 + \left(-s_1\dot{\theta}_1 - s_{12}(\dot{\theta}_1 + \dot{\theta}_2)\right)^2\right)$$

$$KE = \frac{1}{2}\left(c_1^2\dot{\theta}_1^2 + c_{12}^2(\dot{\theta}_1 + \dot{\theta}_2)^2 + s_1^2\dot{\theta}_1^2 + s_{12}^2(\dot{\theta}_1 + \dot{\theta}_2)^2\right)$$

$$c_1c_{12}\dot{\theta}_1(\dot{\theta}_1 + \dot{\theta}_2) + s_1s_{12}\dot{\theta}_1(\dot{\theta}_1 + \dot{\theta}_2)$$

$$KE = \frac{1}{2}\left(\dot{\theta}_1^2 + (\dot{\theta}_1 + \dot{\theta}_2)^2\right) + c_1c_{12}\dot{\theta}_1(\dot{\theta}_1 + \dot{\theta}_2) + s_1s_{12}\dot{\theta}_1(\dot{\theta}_1 + \dot{\theta}_2)$$

From trigonometry

$$\sin(a)\sin(b) + \cos(a)\cos(b) = \cos(a - b)$$

resulting in

$$KE = \frac{1}{2}\left(\dot{\theta}_1^2 + (\dot{\theta}_1 + \dot{\theta}_2)^2\right) + c_2\dot{\theta}_1(\dot{\theta}_1 + \dot{\theta}_2)$$

$$KE = (1 - c_2)\dot{\theta}_1^2 + \frac{1}{2}\dot{\theta}_2^2 + (1 + c_2)\dot{\theta}_1\dot{\theta}_2$$

$$PE = m_2gy_2$$

$$PE = g(c_1 + c_{12})$$

Step 3: Form the LaGrangian:

$$L = KE - PE$$

$$L = \left( (1 + c_2) \dot{\theta}_1^2 + \frac{1}{2} \dot{\theta}_2^2 + (1 + c_2) \dot{\theta}_1 \dot{\theta}_2 \right) - (gc_1 + g(c_1 + c_{12}))$$

$$L = (1 + c_2) \dot{\theta}_1^2 + \frac{1}{2} \dot{\theta}_2^2 + (1 + c_2) \dot{\theta}_1 \dot{\theta}_2 - 2gc_1 - gc_{12}$$

Step 4: Take the partials

$$T_1 = \frac{d}{dt} \left( \frac{\partial L}{\partial \dot{\theta}_1} \right) - \left( \frac{\partial L}{\partial \theta_1} \right)$$

$$T_2 = \frac{d}{dt} \left( \frac{\partial L}{\partial \dot{\theta}_2} \right) - \left( \frac{\partial L}{\partial \theta_2} \right)$$

Starting with  $\theta_1$ :

$$\frac{\partial L}{\partial \dot{\theta}_1} = 2(1 + c_2) \dot{\theta}_1 + (1 + c_2) \dot{\theta}_2$$

$$\frac{d}{dt} \left( \frac{\partial L}{\partial \dot{\theta}_1} \right) = 2(1 + c_2) \ddot{\theta}_1 + (1 + c_2) \ddot{\theta}_2 - 2s_2 \dot{\theta}_1 \dot{\theta}_2 - s_2 \dot{\theta}_2^2$$

$$\frac{\partial L}{\partial \theta_1} = 2gs_1 + gs_{12}$$

Putting it together:

$$T_1 = \frac{d}{dt} \left( \frac{\partial L}{\partial \dot{\theta}_1} \right) - \left( \frac{\partial L}{\partial \theta_1} \right)$$

$$T_1 = 2(1 + c_2) \ddot{\theta}_1 + (1 + c_2) \ddot{\theta}_2 - 2s_2 \dot{\theta}_1 \dot{\theta}_2 - s_2 \dot{\theta}_2^2 - (2gs_1 + gs_{12})$$

Correct Equations from John Craig

$$T_1 = 2(1 + c_2) \ddot{\theta}_1 + (1 + c_2) \ddot{\theta}_2 - 2s_2 \dot{\theta}_1 \dot{\theta}_2 - s_2 \dot{\theta}_2^2 - (2gc_1 + gc_{12})$$

Moving on to  $\theta_2$

$$L = (1 + c_2)\dot{\theta}_1^2 + \frac{1}{2}\dot{\theta}_2^2 + (1 + c_2)\dot{\theta}_1\dot{\theta}_2 - 2gc_1 - gc_{12}$$

Take the partial derivative with respect to  $d\theta_2$ :

$$\frac{\partial L}{\partial \dot{\theta}_2} = \dot{\theta}_2 + (1 + c_2)\dot{\theta}_1$$

Taking the full derivative:

$$\frac{d}{dt} \left( \frac{\partial L}{\partial \dot{\theta}_2} \right) = \ddot{\theta}_2 + (1 + c_2)\ddot{\theta}_1 - s_2\dot{\theta}_1\dot{\theta}_2$$

Now, on the partial with respect to  $q_2$ :

$$\frac{\partial L}{\partial \theta_2} = -s_2\dot{\theta}_1^2 - s_2\dot{\theta}_1\dot{\theta}_2 + gs_{12}$$

Putting it together:

$$T_2 = \frac{d}{dt} \left( \frac{\partial L}{\partial \dot{\theta}_2} \right) - \left( \frac{\partial L}{\partial \theta_2} \right)$$

$$T_2 = \left( \ddot{\theta}_2 + (1 + c_2)\ddot{\theta}_1 - s_2\dot{\theta}_1\dot{\theta}_2 \right) - \left( -s_2\dot{\theta}_1^2 - s_2\dot{\theta}_1\dot{\theta}_2 + gs_{12} \right)$$

$$T_2 = \ddot{\theta}_2 + (1 + c_2)\ddot{\theta}_1 + s_2\dot{\theta}_1^2 - gs_{12}$$

Correct Equations from John Craig:

$$T_2 = \ddot{\theta}_2 + (1 + c_2)\ddot{\theta}_1 + s_2\dot{\theta}_1^2 + gc_{12}$$

Net result:

$$T_1 = 2(1 + c_2)\ddot{\theta}_1 + (1 + c_2)\ddot{\theta}_2 - 2s_2\dot{\theta}_1\dot{\theta}_2 - s_2\dot{\theta}_2^2 + 2gc_1 + gc_{12}$$

$$T_2 = \ddot{\theta}_2 + (1 + c_2)\ddot{\theta}_1 + s_2\dot{\theta}_1^2 + gc_{12}$$

To solve, rewrite this in matrix form:

$$\begin{bmatrix} 2(1 + c_2) & (1 + c_2) \\ (1 + c_2) & 1 \end{bmatrix} \begin{bmatrix} \ddot{\theta}_1 \\ \ddot{\theta}_2 \end{bmatrix} = \begin{bmatrix} T_1 \\ T_2 \end{bmatrix} + \begin{bmatrix} 2s_2\dot{\theta}_1\dot{\theta}_2 + s_2\dot{\theta}_2^2 \\ -s_2\dot{\theta}_1^2 \end{bmatrix} - g \begin{bmatrix} 2c_1 + c_{12} \\ c_{12} \end{bmatrix}$$

Mass Matrix \* Acceleration = Torque - Coriolis Forces + Gravity

Given the joint angles, velocities, gravity, and input torques, you can compute the joint accelerations as

$$\begin{bmatrix} \ddot{\theta}_1 \\ \ddot{\theta}_2 \end{bmatrix} = \begin{bmatrix} 2(1-c_2) & (1-c_2) \\ (1-c_2) & 1 \end{bmatrix}^{-1} \left( \begin{bmatrix} T_1 \\ T_2 \end{bmatrix} - \begin{bmatrix} 2s_2\dot{\theta}_1\dot{\theta}_2 + s_2\dot{\theta}_2^2 \\ -s_2\dot{\theta}_1^2 \end{bmatrix} + \begin{bmatrix} 2gs_1 + gs_{12} \\ gs_{12} \end{bmatrix} \right)$$

Note that the mass matrix is zero when  $\theta_2$  is 180 degrees. This suggests that the acceleration goes to infinity at this position (and should be avoided)

**MatLab Code:**

```
function [ ddQ ] = TwoLinkDynamics( Q, dQ, T )

q1 = Q(1);
q2 = Q(2);

dq1 = dQ(1);
dq2 = dQ(2);

c1 = cos(q1);
s1 = sin(q1);
s2 = sin(q2);
c2 = cos(q2);
s12 = sin(q1+q2);
c12 = cos(q1+q2);

g = 9.8;

M = [ 2*(1+c2) + 0.002,      1+c2 ;
       1+c2           ,      1+0.002 ];

C = [ 2*s2*dq1*dq2 + s2*dq2*dq2 ;
       -s2*dq1*dq1 ];
G = [ 2*g*s1 + g*s12 ;
       g*s12 ];
ddQ = inv(M) * ( T + C + G );

end
```

**Note:** The mass matrix becomes singular at 0 and 180 degrees. To avoid this singularity, 0.01 is added to the diagonal (the arm isn't exactly a point mass)




Image of 2-link arm in free-fall

---

Main Calling Routine:

```
Q = [2; 0.2];
dQ = [0; 0];

T = [0; 0];

t = 0;
dt = 0.01;

while(t < 100)

    ddQ = TwoLinkDynamics(Q, dQ, T);

    dQ = dQ + ddQ * dt;
    Q = Q + dQ*dt;
    t = t + dt;

    x0 = 0;
    y0 = 0;

    x1 = -sin(Q(1));
    y1 = cos(Q(1));

    x2 = x1 - sin(Q(1) + Q(2));
    y2 = y1 + cos(Q(1) + Q(2));

    clf;
    plot([-2,2],[-2,2], 'x');
    hold on;
    plot([-2,2],[0,0], 'g');
    plot([0,0],[-2,2], 'g');

    plot([x0, x1, x2], [y0, y1, y2], 'b.-');
    pause(0.01);

end
```